

# Radiology

## Can Lung US Help Critical Care Clinicians in the Early Diagnosis of Novel Coronavirus (COVID-19) Pneumonia?

Erika Poggiali, Alessandro Dacrema, Davide Bastoni, Valentina Tinelli, Elena Demichele, Pau Mateo Ramos, Teodoro Marcianò, Matteo Silva, Andrea Vercelli, and Andrea Magnacavallo

All authors: Department of Emergency Medicine, Ospedale Guglielmo da Saliceto, via Taverna 49, Piacenza 29121, Italy

Dr. Erika Poggiali

email: [poggiali.erika@gmail.com](mailto:poggiali.erika@gmail.com), [E.Poggiali@ausl.pc.it](mailto:E.Poggiali@ausl.pc.it)

Manuscript Type: Letter to the Editor

Dear Editor:

As reported by Feng et al, chest CT has a pivotal role for the diagnosis and assessment of lung involvement in COVID-19 pneumonia (1). Nowadays CT protocols are used to estimate the pulmonary damage (1,2,3). Unfortunately, CT scanning is not available in all emergency departments. Lung US is a surface imaging technique greatly developed in the last decades and strongly recommended for acute respiratory failure (4). It is commonly used in the emergency department at the bedside for early diagnosis of pneumonia. It is a highly sensitive and specific technique considered as an alternative to chest radiography or CT scanning (5,6). We evaluated the role of lung US in patients who presented to our emergency department with COVID-19 pneumonia. Twelve patients (9 male and 3 female, mean age

63±13 years) with flu-like symptoms in the last 4–10 days and COVID-19 infection underwent bedside lung US and CT. Two patients had emphysema but without need of oxygen therapy at home. None of the patients had severe respiratory distress ( $\text{PaO}_2/\text{FiO}_2$  257–376 mmHg).

In all the patients, we found a diffuse B-pattern with spared areas. Only three patients had posterior subpleural consolidations. Chest CT scan was performed in all 12 patients and showed a strong correlation with US: bilateral lung involvement with ground-glass opacity; five of 12 patients had a crazy-paving pattern. Organizing pneumonia was confirmed in four patients as well as detected with lung US.

We are aware that our data are preliminary and further studies are necessary to confirm the role of lung US in the diagnosis and management of COVID-19 pneumonia, but we strongly recommend the use of bedside US for the early diagnosis of COVID-19 pneumonia in all the patients who presented to the emergency department with flu-like symptoms in novel COVID-19 era.

**Acknowledgments:** The authors would like to thank all the emergency doctors and nurses of their hospital for the help and energy to face such a difficult public health crisis.

## References

1. Feng P, Tianhe Y, Peng S, Shan G, Bo L, Lingli L, Dandan Z, Jiazheng W, Richard L. H, Lian Y, Chuansheng Z. The course of lung changes on chest CT during recovery from 2019 novel coronavirus (COVID-19) Pneumonia. Radiology 2020. doi: 10.1148/radiol.2020200370
2. Chung M, Bernheim A, Mei X, Zhang N, Huang M, Zeng X, Cui J, Xu W, Yang Y, Fayad ZA, Jacobi A, Li K, Li S, Shan H. CT Imaging Features of 2019 Novel Coronavirus (2019-nCoV). Radiology 2020. doi: 10.1148/radiol.2020200230

3. Ai T, Yang Z, Hou H, Zhan C, Chen C, Lv W, Tao Q, Sun Z, Xia L. Correlation of Chest CT and RT-PCR Testing in Coronavirus Disease 2019 (COVID-19) in China: A Report of 1014 Cases. *Radiology* 2020. doi: 10.1148/radiol.2020200642
4. Mongodi S, Pozzi M, Orlando A, Bouhemad B, Stella A, Tavazzi G, Via G, Iotti GA, Mojoli F. Lung ultrasound for daily monitoring of ARDS patients on extracorporeal membrane oxygenation: preliminary experience. *Intensive Care Med.* 2018. doi: 10.1007/s00134-017-4941-7. Epub 2017 Sep 22.
5. Laursen CB, Sloth E, Lassen AT, Christensen Rd, Lambrechtsen J, Madsen PH, Henriksen DP, Davidsen JR, Rasmussen F. Point-of-care ultrasonography in patients admitted with respiratory symptoms: a single-blind, randomised controlled trial. *Lancet Respir Med.* 2014. doi: 10.1016/S2213-2600(14)70135-3
6. Zhou B, Bartholmai BJ, Kalra S, Osborn TG, Zhang X. Lung US Surface Wave Elastography in Interstitial Lung Disease Staging. *Radiology* 2019. doi: 10.1148/radiol.2019181729